



Fecha de presentación: 14/12/2021

Fecha de aceptación: 11/01/2021

Fecha de publicación: 17/02/2021

¿Cómo citar este artículo?

Guardia Puebla, Y., Olivera López, Y., Arias Cedeño, Q., Llanes Cedeño, E., & Eichler Löbermann, B. (enero-abril, 2021). Effect of two chemical pretreatments in the anaerobic hydrolysis of the rice straw. Revista *Márgenes*, 9(1), 20-28. Recuperado de <http://revistas.uniss.edu.cu/index.php/margenes/issue/view/1182>

TITLE: EFFECT OF TWO CHEMICAL PRETREATMENTS IN THE ANAEROBIC HYDROLYSIS OF THE RICE STRAW

TÍTULO: EFECTO DE DOS PRETRATAMIENTOS ALCALINOS EN LA HIDROLISIS ANAEROBIA DE LA PAJA DE ARROZ

Autores: Dr. C Yans Guardia-Puebla¹, Ing. Yasbel Olivera-López², Dr. C Quirino Arias-Cedeño³, Dr. C Edilberto Llanes-Cedeño⁴, Dr. C Bettina Eichler-Löbermann⁵

¹ Ingeniero Hidráulico. Máster en Biotecnología, Mención Ambiental. Doctor en Ciencias Técnicas Agropecuarias. Profesor Titular. Universidad de Granma, Centro de Estudios de Química Aplicada (CEQA), Granma, Cuba. Línea de investigación: Bioenergía y tratamiento de residuos orgánicos. Correo electrónico: yguardiap@udg.co.cu ORCID: <https://orcid.org/0000-0002-1347-0963>

² Ingeniero Químico. Universidad de Granma, Centro de Estudios de Química Aplicada (CEQA), Granma, Cuba. Línea de investigación: Bioenergía y tratamiento de residuos orgánicos. Correo electrónico: yasbel.olivera@nauta.cu ORCID: <https://orcid.org/0000-0001-9081-0725>

³ Licenciado en Química. Doctor en Ciencias Químicas. Profesor Titular. Universidad de Granma, Centro de Estudios de Química Aplicada (CEQA), Granma, Cuba. Línea de investigación: Síntesis orgánica. Correo electrónico: gariasc@udg.co.cu ORCID: <https://orcid.org/0000-0001-6024-9877>

⁴ Ingeniero Mecánico Automotriz. Máster en Eficiencia Energética. Doctor en Ingeniería Rural. Profesor Titular. Universidad Internacional SEK Ecuador, Facultad de Arquitectura e Ingenierías, Quito, Ecuador. Línea de investigación: Eficiencia energética y energía. Correo electrónico: antonio.llanes@uisek.edu.ec ORCID: <https://orcid.org/0000-0001-6739-7661>

⁵ Prof. Dr. Agr. Habil., Universidad de Rostock, Departamento de Ciencias de las Plantas, Facultad de Agricultura, Rostock, Alemania. Línea de investigación: Ciencias de las Plantas. Correo electrónico: bettina.eichler@uni-rostock.de ORCID: <https://orcid.org/0000-0001-8306-0452>

* This study was partly supported by the Federal Ministry of Education and Research (BMBF) project InFertRes (CUB17WTZ-042).

Márgenes publica sus artículos bajo una [Licencia Creative Commons Atribución-NoComercial-SinDerivar 4.0 Internacional](https://creativecommons.org/licenses/by-nc/4.0/)



<http://revistas.uniss.edu.cu/index.php/margenes>
margenes@uniss.edu.cu

ABSTRACT

Rice (*Oryza sativa*) is an important staple food for approximately half of the world population. In addition to the grains, also the straw can be used if not needed as organic fertilizer for the fields. Although rice straw has potential to be used for anaerobic digestion (AD) to produce biogas, its large-scale application is still limited. The utilization of rice straw for AD must particularly be developed regarding the pretreatment of the lignin biomass for the hydrolysis process and the biogas yield and quality. Among the pretreatment methods, the alkalization method was found to be effective. In the present research the effects of rice straw pretreatment were analyzed following a multifactorial approach. For this purpose, calcium hydroxide (Ca(OH)_2) and potassium hydroxide (KOH) were applied at different concentrations. Furthermore, the reaction time and inoculum-to-substrate ratio were included as part of the study. The highest hydrolysis yield was observed at a reaction time of 4 h, with an alkali concentration of 10 g per g rice dry matter, and an inoculum-to-substrate ratio of 50%. Here also the maximum concentration of volatile fatty acids was observed with $187 \pm 30 \text{ mg L}^{-1}$. The results showed that the additional effort of a pretreatment of rice straw can be worthwhile to improve the AD and to increase the biogas yield.

Keywords: chemical pretreatment; anaerobic digestion; rice straw; hydrolysis.

RESUMEN

El arroz (*Oryza sativa*) es un alimento básico importante para aproximadamente la mitad de la población mundial. Además de los granos, la paja de arroz también se puede usar si no se necesita como fertilizante orgánico para los campos. Aunque la paja de arroz tiene el potencial de utilizarse para la digestión anaerobia (DA) para producir biogás, su aplicación a gran escala aún es limitada. La utilización de la paja de arroz para la DA debe desarrollarse particularmente en lo que respecta al pretratamiento de la lignina para el proceso de hidrólisis y el rendimiento y la calidad del biogás. Entre los métodos de pretratamiento, se ha comprobado que la alcalinización es la más efectiva. En el presente estudio, se investigaron los efectos del pretratamiento

Márgenes publica sus artículos bajo una [Licencia Creative Commons Atribución-NoComercial-SinDerivar 4.0 Internacional](http://creativecommons.org/licenses/by-nc/4.0/)



<http://revistas.uniss.edu.cu/index.php/margenes>
margenes@uniss.edu.cu

de la paja de arroz en un enfoque multifactorial. Para ello se aplicaron el hidróxido de calcio ($\text{Ca}(\text{OH})_2$) y el hidróxido de potasio (KOH) a diferentes concentraciones. Además, el tiempo de reacción y la relación de inóculo/sustrato se incluyeron en la investigación. El mayor rendimiento de la hidrólisis se observó a un tiempo de reacción de 4 h, con una concentración de álcali de 10 g/g de materia seca y una relación de inóculo sustrato del 50 %. También, la máxima concentración de los ácidos grasos volátiles (AGV) alcanzada fue de $187 \pm 30 \text{ mg L}^{-1}$. Los resultados mostraron que el pretratamiento de la paja de arroz vale la pena para mejorar la DA y aumentar el rendimiento del biogás.

Palabras clave: pretratamiento químico; digestión anaerobia; paja de arroz; hidrólisis.

INTRODUCTION

Rice straw is one of the most abundant renewable lignocelluloses crop residues (LCR) with the most availability in the world. Rice Straw is commonly used for animal feed and as a fuel for cooking and heating homes. However, the largest amounts of rice straw remain unused in the field and burned in the open fields causing serious environmental problems (Guan et al., 2018). Anaerobic digestion (AD) to produce biogas can offer promising benefits for using rice straw and mitigating air pollution. It has been confirmed that AD is an attractive technology for simultaneous clean bioenergy production and waste treatment. Biogas and digestate are the two beneficial products of AD. Digestate is rich in nitrogen and organic matter and can be used in agriculture as a biofertilizer or soil improver (Carrere et al., 2016). However, the inherent characteristics of rice straw made it resistant to enzymatic degradation by anaerobic microbes. For that reason, pretreatment of rice straw prior to AD process has been proven to be necessary to improve biodegradability and biogas production. Chemical pretreatment has been defined as the most promising method to improve the bioconversion of cellulose in order to improve the enzymatic accessibility and thus, facilitate the subsequent anaerobic treatment (Sabeeh, Zeshan, Liaquat, & Maryam, 2020). Various pretreatment protocols have been used for the biomethanation of rice straw involving heating and the use of various amounts of chemicals (Boonsombuti, Trisinsub, & Luengnaruemitchai, 2019;

Márgenes publica sus artículos bajo una [Licencia Creative Commons Atribución-NoComercial-SinDerivar 4.0 Internacional](https://creativecommons.org/licenses/by-nc/4.0/)



<http://revistas.uniss.edu.cu/index.php/margenes>
margenes@uniss.edu.cu

Dong, Cao, Zhao, Liu, & Ren, 2018; Kim, Kim, Nam, & Choi, 2018; Mancini, Papirio, Riccardelli, Lens, & Esposito, 2017; Wua et al., 2017) However, alkali pretreatments are commonly investigated at elevated temperature (over 100°C) with the drawbacks of special equipment requirement, higher treatment costs and high energy consumption (Du, Qian, Xi, & Lü, 2019). Hence, the major objective of the present research was to minimize the alkali requirement at low temperature during pretreatment to enhance the hydrolysis of rice straw. In the present research the effects of rice straw pretreatment were analyzed in a multifactorial approach. For this purpose, calcium hydroxide ($\text{Ca}(\text{OH})_2$) and potassium hydroxide (KOH) were applied at different concentrations.

MATERIAL AND METHODS

Rice straw was air-dried and milled using hammer mill grinder and sieved to select straw of desired particle size. The cellulose, hemicellulose and acid-insoluble lignin content in the rice straw sample was determined by the Van Soest method (Van Soest, Robertson, & Lewis, 1991). The detection of total solids (TS) and volatile solids (VS) was carried out in accordance with the standard methods of APHA (APHA 1995). Slurry obtained from a cow dung biometanation industrial plant operated at 37 °C was used as an inoculum for hydrolysis from rice straw. The pH of the inoculum was 7. To study the best conditions of hydrolysis of rice straw residues, a factorial experimental design was used. Four experimental factors were considered: type of alkali compound ($\text{Ca}(\text{OH})_2$, KOH), time of reaction (2, 3, and 4 h), concentration of alkali compound (4, 7, and 10 g/g dry matter), and inoculum-to-substrate ratio (20, 35, and 50 %) (Figure 1).






Figure 1. Schematic performance of the experiment

Source: Own elaboration

In the designed tests, 4 g of rice straw was added to 100 mL of each alkali solution during the corresponding pretreatment time and kept at 35 °C with circulating water. Then, the alkali solution was drained and the corresponding inoculum-to substrate ratio was added to each bottle and subsequently made up to volume with water to a volume of 100 mL. All tests were performed at 35 °C using 250 mL borosilicate glass reactors with 100 mL working volume, at an initial pH of 7.0. An untreated sample of rice straw was simultaneously tested as the control. Each experiment was performed for 10 days. VFA samples were prepared by centrifuging at 13000 rpm for 5 min before filtering through a 0.2 μm nylon membrane. VFA concentrations were measured using a gas chromatograph (GC-2014, Shimadzu, Japan) equipped with a flame ionization detector (FID).

RESULTS AND DISCUSSION

Composition of rice straw

Rice straw, like many lignocellulosic biomasses, possess several properties that makes them suitable as feedstocks for bio-chemical conversion. Table 1 shows the chemical characteristics of the rice straw used in this study. The results showed that the rice straw was composed of 48.7% cellulose, 22.2% hemicellulose, and 3.1% lignin. Thus, the rice

Márgenes publica sus artículos bajo una [Licencia Creative Commons Atribución-NoComercial-SinDerivar 4.0 Internacional](http://creativecommons.org/licenses/by-nc/4.0/)



<http://revistas.uniss.edu.cu/index.php/margenes>
margenes@uniss.edu.cu

straw was considered as a promising carbon source for microbial fermentation. On the other hand, rice straw contained 90.3% (w/w) total solid; meanwhile, volatile solids constituted 77.5% of the TS.


Table 1. Characteristics of rice straw

Celulose (%)	Hemicelulose (%)	Lignin (%)	pH	TS (%)	VS (% of TS)	Na (mg/L)	K (mg/L)
48.7	22.2	3.1	6.42	90.3	82.1	10	150
P (mg/L)	Ca (mg/L)	Cd (mg/L)	Cr (mg/L)	Cu (mg/L)	Mg (mg/L)	Ni (mg/L)	Pb (mg/L)
37.5	35.9	0.017	0.023	0.053	17.6	0.047	1.06

Source: Own elaboration

Effect of pretreatments on hydrolysis process

Alkali pretreatment is capable of improving rice straw biodegradability by releasing organic soluble fraction. A fixed quantity of rice straw was exposed to various concentrations of $\text{Ca}(\text{OH})_2$ and KOH for varying duration to optimize the pretreatment at low temperature. The efficacy of the pretreatment on the hydrolysis stage was monitored in terms of VFA production. The VFA concentrations of the pretreated rice straw is shown in Figure 2.



Márgenes publica sus artículos bajo una [Licencia Creative Commons Atribución-NoComercial-SinDerivar 4.0 Internacional](https://creativecommons.org/licenses/by-nc/4.0/)



<http://revistas.uniss.edu.cu/index.php/margenes>
margenes@uniss.edu.cu




Figure 2. VFA concentration of pretreated rice straw; a) and b) acetic acid; c and d) butyric acid; and e) and f) cumulated total VFA.

Source: Own elaboration

VFA in the anaerobic reactors mainly comprised acetic and propionic acids. No significant accumulation of butyric and iso-butyric acids was observed, probably because of the sufficient butyric-degrading syntrophs in the inoculum, which rapidly converted these acids to acetic acid. Total VFA production increased from 89 ± 12 mg/L for untreated rice straw to 187 ± 30 mg/L when rice straw was pretreated for 4 hours with 10 g per g rice dry matter of $\text{Ca}(\text{OH})_2$ and used an inoculum-to-substrate ratio of 50 %. Compared to KOH, the $\text{Ca}(\text{OH})_2$ had more significant influence on the VFA production. ANOVA confirmed that each concentration of alkali compounds significantly ($p < 0.05$) influenced the production of VFA. Economically, the optimal amount of $\text{Ca}(\text{OH})_2$ for the

pretreatment of rice straw was considered 4 g per g rice dry matter due to the observation of not significant differences at dose of 10 g per g rice dry matter. Hence, for further research, rice straw will be pretreated with 5% (g/gTS) $\text{Ca}(\text{OH})_2$ for 4 hours at 35 °C temperature and evaluated for biomethanation. The preliminary results suggested that from the same amount of raw material loaded, the pretreated rice straw with $\text{Ca}(\text{OH})_2$ the hydrolysis yield is highest compared to untreated rice straw.

CONCLUSIONS

The results of this study show that $\text{Ca}(\text{OH})_2$ pretreatment could significantly enhance the hydrolysis of milled rice straw. Maximum production of VFA (187 mg/L) was achieved with rice straw treated with 10% $\text{Ca}(\text{OH})_2$ at 35 °C temperature for 4 hours. Alkali pretreatment increased the production of VFA by over 47%, obtained from milled rice straw without alkali treatment. $\text{Ca}(\text{OH})_2$ pretreatment is a promising pretreatment process that can be carried out at low temperature.

BIBLIOGRAPHIC REFERENCES

- Boonsombuti, A., Trisinsub, O., & Luengnaruemitchai, A. (2019). Comparative study of three chemical pretreatments and their effects on the structural changes of rice straw and butanol production. *Waste and Biomass Valorization*. doi:10.1007/s12649-019-00622-z
- Carrere, H., Antonopoulou, G., Affes, R., Passos, F., Battimelli, A., Lyberatos, G., & Ferrer, I. (2016). Review of feedstock pretreatment strategies for improved anaerobic digestion: From lab-scale research to full-scale application. *Bioresource Technology*, 199, 386-397. doi:10.1016/j.biortech.2015.09.007
- Dong, L., Cao, G., Zhao, L., Liu, B., & Ren, N. (2018). Alkali/Urea pretreatment of rice straw at low temperature for enhanced biological hydrogen production. *Bioresource Technology*. doi:10.1016/j.biortech.2018.05.055
- Du, J., Qian, Y., Xi, Y., & Lü, X. (2019). Hydrothermal and alkaline thermal pretreatment at mild temperature in solid state for physicochemical properties and biogas production from anaerobic digestion of rice straw. *Renewable Energy*, 139, 261-267. doi:10.1016/j.renene.2019.01.097
- Guan, R., Li, X., Wachemo, A., Yuana, H., Liu, Y., Zou, D., & Gao, J. (2018). Enhancing anaerobic digestion performance and degradation of lignocellulosic components of rice straw by combined biological and chemical pretreatment. *Science of the Total Environment*, 637-638, 9-17. doi:10.1016/j.scitotenv.2018.04.366



-
- Kim, M., Kim, B.-C., Nam, K., & Choi, Y. (2018). Effect of pretreatment solutions and conditions on decomposition and anaerobic digestion of lignocellulosic biomass in rice straw. *Biochemical Engineering Journal*. doi:10.1016/j.bej.2018.09.012
- Mancini, G., Papirio, S., Riccardelli, G., Lens, P., & Esposito, G. (2017). Trace elements dosing and alkaline pretreatment in the anaerobic digestion of rice straw. *Bioresource Technology*. doi:10.1016/j.biortech.2017.10.001
- Sabeeh, M., Zeshan, Liaquat, R., & Maryam, A. (2020). Effect of alkaline and alkaline-photocatalytic pretreatment on characteristics and biogas production of rice straw. *Bioresource Technology*, 309, 123449. doi:10.1016/j.biortech.2020.123449
- Van Soest, P. J., Robertson, J. B., & Lewis, B. A. (1991). Methods for dietary fiber, neutral detergent fiber, and nonstarch polysaccharides in relation to animal nutrition. *J Dairy Sci*, 74(10), 3583-3597. doi:10.3168/jds.S0022-0302(91)78551-2
- Wua, H., Dai, X., Zhou, S.-L., Gan, Y.-Y., Xiong, Z.-Y., Qin, Y.-H., & Wang, C.-W. (2017). Ultrasound-assisted alkaline pretreatment for enhancing the enzymatic hydrolysis of rice straw by using the heat energy dissipated from ultrasonication. *Bioresource Technology*, 241(70-74). doi:10.1016/j.biortech.2017.05.090

